Nepenthes of Gunung Ulu Kali

Roger G. Shivas

Botany Department, National University of Malaysia, Bangi, Selangor, Malaysia

The Genting Highlands are situated about 40 km northeast of Kuala Lumpur in tropical Malaysia. The highest mountain in this part of the range is Gunung Ulu Kali (1800 m). Presently the area is being developed as a tourist resort and boasts four hotels, a golf course, and a casino.

Gunung Ulu Kali is often shrouded in cloud and rainfall in the area is high. The temperature ranges between the extremes of hot days (30° C) and cool nights (10° C).

The vegetation on the ridges and crests is upper montane moss forest characterized by stunted trees of *Dacrydium* and *Leptospermum*, several species of *Rhododendron* and other cricaceous shrubs and many epiphytic orchids. The top soil is peat, which is often covered in dense mats of sphagnum moss.

Three species of Nepenthes have been recorded from Gunung Ulu Kali, Nepenthes macfarlanei, Nepenthes gracillima var major and Nepenthes sanguinea (Stone, 1981). During 1982 and 1983, I made several trips to Gunung Ulu Kali to observe and collect Nepenthes.

Nepenthes macfarlanei is a robust plant with waxy leaves and was the most abundant plant species in the moss forest. This species was readily distinguished by its cream-coloured, crimson-spotted upper pitchers and large, ovate (10 to 15 cm high and 5 to 8 cm wide), mottled red lower pitchers. Plants often reached lengths greater than 4 m.

Nepenthes gracillima, although not as plentiful as Nepenthes macfarlanei, was easily identified by its dark purplishblack, cylindrical pitchers (10 to 20 cm high and 1 to 3 cm wide). Mature plants were vines greater than 5 m in length. The form of this species found on Gunung Ulu Kali most closely fits Nepenthes gracillima var major (Ridley, 1924).


Nepenthes sanguinea was rarely found on the ridges and crests of Gunung Ulu Kali. However, this species was found along roadside embankments at an altitude 300 m lower than the summit. *Nepenthes sanguinea* was conspicuous with its wide peristome (1 to 2 cm) and cylindrical pitchers (10 to 20 cm high and 3 to 6 cm wide). The colour of the pitchers varied from pale red and green to a dark, uniform blood red.

Literature Cited

Ridley, H.N. (1924) The Flora of the Malay Peninsula, III. London, L. Reeve & Co., p. 22.

Stone, B.C. (1981) The summit flora of Gunung Ulu Kali (Pahang, Malaysia). Federation Museums Journal, Kuala Lumpur, Volume 26 (Part I), N.S.

(See front and back covers as well as photos, pages 66-7).



Drawing by Ron Fleming



Upper pitcher of Nepenthes macfarlanei.

Photo by Roger Shivas



Lower pitcher of Nepenthes macfarlanei.



Figure 1—The smaller plants are green S. $alata \times minor$ (North Carolina), and the larger plants are the red-throated S. $alata \times S$. minor (Okefenokee).



Figure 2—S. alata red-throated \times S. flava heavy-veined.

Photos above by Steve Clemesha.



Distinctive purplish-black pitchers of Nepenthes gracillima.



CARNIVOROUS PLANT NEWSLETTER

Official Journal of the International Carnivorous Plant Society

Volume 12, Number 3 September 1983

COVER PHOTOS

Front: Rosette plant and lower pitcher of Nepenthes sanguinea.

Back: Lower pitchers of Nepenthes sanguinea.

See story, page 65.

Photos by Roger Shivas

The co-editors of CPN would like everyone to pay particular attention to the following policies regarding your dues to the ICPS.

All correspondence regarding dues, address changes and missing issues should be sent to Mrs. Pat Hansen, 3321 Hamell Rd., Fullerton, CA 92635. DO NOT SEND TO THE CO-EDITORS. Checks for subscriptions and reprints should be made payable to CSUF FOUNDATION-ICPS.

All material for publication, comments and general correspondence about your plants, field trips or special noteworthy events relating to CP should be directed to one of the co-editors. We are interested in all news related to carnivorous plants and rely on the membership to supply us with this information so that we can share it with others.

Views expressed in this publication are those of the authors, not necessarily the editorial staff. Copy deadline for the December issue is October 15, 1983.

CO-EDITORS:

D. E. Schnell, Rt. 1, Box 145C, Pulaski, VA 24301

J. A. Mazrimas, 329 Helen Way, Livermore, CA 94550

T. L. Mellichamp, Dept. of Biology, UNCC, Charlotte, NC 28223

Leo Song, Dept. of Biology, California State University, Fullerton, CA 92634

Seed Bank: Patrick Dwyer, St. Michael's Episcopal Church, 49 Killean Park, Albany, NY 12205, USA.

BUSINESS MANAGER: Mrs. Pat Hansen, c/o The Fullerton Arboretum

PUBLISHER: The International Carnivorous Plant Society by the Fullerton Arboretum,
California State University Fullerton, CA 92634, Published quarterly

California State University, Fullerton, CA 92634. Published quarterly with one volume annually. Printer: Kandid Litho, 129 Agostino Rd., San Gabriel, CA 91776. Circulation: 656 (160 new, 496 renewal). Dues: \$10.00 annually, \$15.00 foreign. Reprints available by volume only. \$1983 Carnivorous Plant Newsletter. All rights reserved.

NEWSLETTER CARNIVOROUS PLANT



